International Symposium dedicated to the 70th anniversary of Byurakan Astrophysical Observatory (Byurakan, 19-23 September 2016)
Non-Stable Universe: Energetic Resources, Activity Phenomena and Evolutionary Processes

The impact of bulges on the radial distribution of supernovae in disc galaxies

A. A. Hakobyan[1*], A. G. Karapetyan[1†], L. V. Barkhudaryan[1], G. A. Mamon[2], D. Kunth[2], A. R. Petrosian[1], V. Adibekyan[3], L. S. Aramyan[1] and M. Turatto[4]

[1] Byurakan Astrophysical Observatory, Byurakan, Armenia
[2] Institut d'Astrophysique de Paris, Paris, France
[3] Instituto de Astrofísica e Ciência do Espaço, Porto, Portugal
[4] Osservatorio Astronomico di Padova, Padova, Italy

arXiv:1609.08319v1 [astro-ph.GA] 27 Sep 2016

Abstract

We present an analysis of the impact of bulges on the radial distributions of the different types of supernovae (SNe) in the stellar discs of host galaxies with various morphologies. We find that in Sa–Sm galaxies, all core-collapse (CC) and vast majority of SNe Ia belong to the disc, rather than the bulge component. The radial distribution of SNe Ia in S0–S0/a galaxies is inconsistent with their distribution in Sa–Sm hosts, which is probably due to the contribution of the outer bulge SNe Ia in S0–S0/a galaxies. The radial distributions of both types of SNe are similar in all the subsamples of Sa–Sbc and Sc–Sm galaxies. These results confirm that the old bulges of Sa–Sm galaxies are not significant producers of Type Ia SNe, while the bulge populations are significant for SNe Ia only in S0–S0/a galaxies.

Summary

This is a brief summary of Hakobyan et al. (2016), written for a short contribution in the symposium *"Non-Stable Universe: Energetic Resources, Activity Phenomena and Evolutionary Processes"* dedicated to the 70th anniversary of Byurakan Astrophysical Observatory. In the mentioned paper, using a well-defined and homogeneous sample of SNe and their host galaxies from the coverage of Sloan Digital Sky Survey (Hakobyan et al., 2012, 2014), we analysed the impact of bulges on the radial distributions of the different types of SNe in host galaxies with various morphologies. Our sample consists of 419 nearby (≤ 100 Mpc), low-inclination ($i \leq 60^\circ$), and morphologically non-disturbed S0–Sm galaxies, hosting 500 SNe in total.

All the results that we summarize below concerning the impact of bulges on the radial distributions of SNe in hosts can be explained considering that the old bulges of Sa–Sm galax-

*E-mail: hakobyan@bao.sci.am
†E-mail: karapetyan@bao.sci.am

ies are not significant producers of Type Ia SNe, while the bulge populations are significant for SNe Ia only in S0–S0/a galaxies.

1. In Sa–Sm galaxies, all CC and the vast majority of Type Ia SNe belong to the disc, rather than the bulge component. The result suggests that the rate of SNe Ia in spiral galaxies is dominated by a relatively young/intermediate progenitor population (e.g. Mannucci et al., 2005; Hakobyan et al., 2011; Li et al., 2011). In addition, this observational fact makes the deprojection of galactocentric radii (Hakobyan et al., 2009) of both types of SNe a key point in the statistical studies of their distributions.

2. The radial distribution of Type Ia SNe in S0–S0/a galaxies is inconsistent with that in Sa–Sm hosts. This inconsistency is mostly attributed to the contribution by SNe Ia in the outer bulges of S0–S0/a galaxies. In these hosts, the relative fraction of bulge to disc SNe Ia is probably changed in comparison with that in Sa–Sm hosts, because the progenitor population from the discs of S0–S0/a galaxies should be much lower due to the lower number of young/intermediate stellar populations.

3. The inner fractions ($R_{\rm SN} \leq 0.3\,R_{25}$, where the bulge stars are tipically located) of Type Ia and CC SNe are not statistically different one from another in morphologically non-disturbed spiral hosts. This result additionally confirms that the old bulges of Sa–Sm galaxies are not significant producers of Type Ia SNe.

Acknowledgments

This work was supported by the RA MES State Committee of Science, in the frames of the research project number 15T–1C129. This work was made possible in part by a research grant from the Armenian National Science and Education Fund (ANSEF) based in New York, USA.

References

Hakobyan, A. A., Adibekyan, V. Z., Aramyan, L. S., Petrosian, A. R., Gomes, J. M., Mamon, G. A., Kunth, D., & Turatto, M. 2012, A&A, 544, A81

Hakobyan, A. A., et al. 2016, MNRAS, 456, 2848

Hakobyan, A. A., Mamon, G. A., Petrosian, A. R., Kunth, D., & Turatto, M. 2009, A&A, 508, 1259

Hakobyan, A. A., et al. 2014, MNRAS, 444, 2428

Hakobyan, A. A., et al. 2011, Astrophysics, 54, 301

Li, W., Chornock, R., Leaman, J., Filippenko, A. V., Poznanski, D., Wang, X., Ganeshalingam, M., & Mannucci, F. 2011, MNRAS, 412, 1473

Mannucci, F., Della Valle, M., Panagia, N., Cappellaro, E., Cresci, G., Maiolino, R., Petrosian, A., & Turatto, M. 2005, A&A, 433, 807